



# **BONE HEALTH REFERENCES**

Updated December 2020

---

Compiled by the Bone Health Special Interest Group

*\* References added in 2020 review are in Blue*

	<u>PAGE</u>
<b>EXERCISE AND BONE HEALTH: REVIEWS AND META-ANALYSES</b>	2
<b>MECHANICAL MODELS RELATED TO FRACTURE RISK AND TO EXERCISE</b>	4
<b>IN VIVO MODELS RELATED TO FRACTURE RISK AND TO EXERCISE</b>	6
<b>PREDICTORS AND EFFECTS OF THORACIC HYPERKYPHOSIS</b>	7
<b>EXERCISE INTERVENTIONS FOR KYPHOSIS AND/OR VERTEBRAL FRACTURE</b>	10
<b>EPIDEMIOLOGY AND RISK OF FRACTURE</b>	13
<b>FLEXION EXERCISES</b>	15
<b>OTHER EXERCISE AND ACTIVITY INTERVENTIONS, ASSESSMENTS, &amp; MEDICAL MANAGEMENT FOR BONE HEALTH</b>	15
<b>HIP FRACTURE</b>	20
<b>VIBRATION</b>	21
<b>MEN</b>	23
<b>CHILDREN/ADOLESCENTS/YOUNG ADULTS</b>	24

## **1. EXERCISE AND BONE HEALTH: REVIEWS AND META-ANALYSES**

Armstrong JJ, Rodrigues IB, Wasiuta T, MacDermid JC. Quality assessment of osteoporosis clinical practice guidelines for physical activity and safe movement: an AGREE II appraisal. *Arch Osteoporos*. 2016;11:6. doi:10.1007/s11657-016-0260-9

Auais MA, Eliayyan O, Mayo NE. Extended exercise rehabilitation after hip fracture improves patients' physical function: a systematic review and meta-analysis. *Phys Ther* 2012;92:1437-1451.

Babatunde O; Forsyth J; Gidlo C. A Meta-analysis of High-impact Exercises for Enhancing Bone Health in Premenopausal Women. *Osteoporos Int* 2012;23:109-19.

Bansal S, Katzman WB, Giangregorio LM. Exercise for Improving Age-Related Hyperkyphotic Posture: A Systematic Review. *Arch Phys Med Rehabil* Jul 2013.

Beck, B. R., et al. (2016). "Exercise and Sports Science Australia (ESSA) position statement on exercise prescription for the prevention and management of osteoporosis." [J Sci Med Sport](#).

de Kam D, Smulders E, Weerdesteyn V, Smits-Engelsman BC. Exercise interventions to reduce fall-related fractures and their risk factors in individuals with low bone density: a systematic review of randomized controlled trials. *Osteoporos Int* 2009 Dec;20(12):2111-25.

Chudyk AM, Jutai JW, Petrella RJ, Speechley M. Systematic review of hip fracture rehabilitation practices in the elderly. *Arch Phys Med Rehabil* 2009;90:246-262.

Dusdal K, Grundmanis, Luttin K, et al. Effects of therapeutic exercise for persons with osteoporotic vertebral fractures: a systematic review. *Osteoporos Int* 2011;22:755-769.

Freehan LM, Beck CA, Harris SR et al. Exercise prescription after fragility fracture in older adults: a scoping review. *Osteoporos Int* 2011;22:1289-1322.

Gibbs JC, MacIntyre NJ, Ponzano M, et al. Exercise for improving outcomes after osteoporotic vertebral fracture. *Cochrane Database Syst Rev*. 2019;7(7):CD008618. Published 2019 Jul 5. doi:10.1002/14651858.CD008618.pub3

Gomez-Bruton A, Gonzalez-Aguero A, Gomez-Cabello et al. Is Bone Tissue Really Affected by Swimming? A systematic review. *PLoS ONE* 2013;8:e70119.

Gomez-Bruton, A., et al. (2017). "Plyometric exercise and bone health in children and adolescents: a systematic review." *World J Pediatr*.

Gomez-Cabello A, Ara I, Gonzalez-Aguero A, Casajus JA, Vicente-Rodriguez G. Effects of training on bone mass in older adults: a systematic review. *Sports Med* 2012;42(4):301-325.

Granacher U, Gollhofer A, Hortobagyi T, Kressig RW, Muehlbauer T. The importance of trunk muscle strength for balance, functional performance, and fall prevention in seniors: a systematic review. *Sports Med* 2013;43(7):627-641.

Gregg EW, Pereira MA, Caspersen CJ. Physical activity, falls, and fractures among older adults: a review of the epidemiologic evidence. *J Am Geriatr Soc* 2000 Aug;48(8):883-93. Review.

Hackney ME, & Wolf SL. Impact of Tai Chi Chu'an practice on balance and mobility in older adults: an integrative review of 20 years of research. *J Geriatr Phys Ther* 2014;37(3), 127-135.

Hagen KB, Dagfinrud H, Moe RH, et al. Exercise therapy for bone and muscle health: an overview of systematic reviews. *BMC Med* 2012;10:167.

Hamilton CJ, Swan VJ, Jamal SA. The effects of exercise and physical activity participation on bone mass and geometry in postmenopausal women: a systematic review of pQCT studies. *Osteoporos Int* 2010 Jan;21(1):11-23.

Handoll HHG, Sherrington C, Mak JCS. Interventions for improving mobility after hip fracture surgery in adults. *Cochrane Database of Systematic Reviews* 2011;issue 3. Art No: CD001704.

Howe TE, Shea B, Dawson LJ, et al. Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Database of Systematic Reviews* 2011;issue 7. Art. No.: CD000333.

Jensen AL, Lomborg K, Wind G, Langdahl BL. Effectiveness and characteristics of multifaceted osteoporosis group education--a systematic review. *Osteoporos Int* 2014;25(4):1209-1224.

Exercise prescription for people with osteoporotic vertebral fracture." *Br J Sports Med* 49(7): 489-490.

Kelley GA, Kelley KS, Kohrt WM. Effects of ground and joint reaction force exercise on lumbar spine and femoral neck bone mineral density in postmenopausal women: a meta-analysis of randomized clinical trials. *BMC Musculoskeletal Disorders* 2012;13:177.

Kemmler W, Haberle L, von Stengel S. Effects of exercise on fracture reduction in older adults: a systematic review and meta-analysis. *Osteoporos Int* 2013;24(7):1937-1950.

Kohrt WM, Bloomfield SA, Little KD, Nelson ME, Yingling VR; American College of Sports Medicine. American College of Sports Medicine Position Stand: physical activity and bone health. *Med Sci Sports Exerc* 2004 Nov;36(11):1985-96. Review.

Kasch R, Merk H, Pinto RZ. "Exercise prescription for people with osteoporotic vertebral fracture." *Br J Sports Med* 2015;49(7):489-490.

Lau RW, Liao LR, Yu F, Teo T, Chung RC, Pang MY. The effects of whole body vibration therapy on bone mineral density and leg muscle strength in older adults: a systematic review and meta-analysis. *Clinical Rehabilitation* Nov 2011;25(11):975-988.

Ma D, Wu L, He Z. Effects of walking on the preservation of bone mineral density in perimenopausal and postmenopausal women: a systematic review and meta-analysis. *Menopause* 2013;20(11):1216-1226.

Marques, E. A., et al. "Exercise effects on bone mineral density in older adults: a meta-analysis of randomized controlled trials." *Age (Dordr)* 2012;34(6):1493-1515.

Martyn-St James M, Carroll S. Effects of different impact exercise modalities on bone mineral density in premenopausal women: a meta-analysis. *J Bone Miner Metab* 2010 May;28(3):251-67.

Martyn-St James M, Carroll S. A meta-analysis of impact exercise on postmenopausal bone loss: the case for mixed loading exercise programmes. *Br J Sports Med* 2009 Dec;43(12):898-908.

Merriman H, Jackson K. The effects of whole-body vibration training in aging adults: a systematic review. *J Geriatr Phys Ther* 2009;32(3):134-145.

Moayyeri A. The association between physical activity and osteoporotic fractures: a review of the evidence and implications for future research. *Ann Epidemiol* 2008 Nov;18(11):827-35.

Nikander R, Sievanen H, Heinonen A, Daly R, Uusi-Rasi K, Kannus, P. Targeted exercise against osteoporosis: A systematic review and meta-analysis for optimising bone strength throughout life. *BMC Medicine* 2010, 8:47.

Olmedillas H; Gonzalez-Aguero A; Vicente-Rodriguez G. Cycling and Bone Health: a systematic review. *BMC Med* 2012;10:168.

Pfeifer M, Sinaki M, Geusens P, Boonen S, Preisinger E, Minne HW; ASBMR Working Group on Musculoskeletal Rehabilitation. Musculoskeletal rehabilitation in osteoporosis: a review. *J Bone Miner Res* 2004 Aug;19(8):1208-14. Review.

Polidoulis I, Beyene J, Cheung AM. The effect of exercise on pQCT parameters of bone structure and strength in postmenopausal women – a systematic review and meta-analysis of randomized controlled trials. *Osteoporos Int* 2012;23:39-51.

Rehn B, Lidstrom J, Skoglund J, Lindstrom B. Effects on leg muscular performance from WBV exercise: a systematic review. *Scand J Med Sci Sports* Feb 2007;17(1):2-11.

Sinaki M, Pfeifer M, Preisinger E, Itoi E, Rizzoli R, Boonen S, Geusens P, Minne HW. The role of exercise in the treatment of osteoporosis. *Curr Osteoporos Rep* 2010 Sep;8(3):138-44.

Slatkowska L, Alibhai SMH, Beyene J, Cheung AM. Effect of whole-body vibration on BMD: a systematic review and meta-analysis. *Osteoporos Int* 2010;21:1969-1980.

Turner CH, Robling AG. Designing exercise regimens to increase bone strength. *Exerc Sports Sci Rev* 2003;31:45-50.

Wilhelm, M, Oskovensky G, Emery K, Manno C, Valek K, Cook C. Effect of resistance exercises on function in older adults with osteoporosis or osteopenia: a systematic review. *Physiother Can* 2012;64(4): 386-394.

## **2. MECHANICAL MODELS RELATED TO FRACTURE RISK AND TO EXERCISE**

Adams MA, Pollintine P, Tobias JH, Wakley GK, Dolan P. Intervertebral disc degeneration can predispose to anterior vertebral fractures in the thoracolumbar spine. *J Bone Miner Res* 2006 Sep;21(9):1409-16.

Bouxsein ML, Melton LJ 3rd, Riggs BL, Muller J, Atkinson EJ, Oberg AL, Robb RA, Camp JJ, Rouleau PA, McCollough CH, Khosla S. Age- and sex-specific differences in the factor of risk for vertebral fracture: a population-based study using QCT. *J Bone Miner Res* 2006 Sep;21(9):1475-82.

Briggs AM, Greig AM, Wark JD, Fazzalari NL, Bennell KL. A review of anatomical and mechanical factors affecting vertebral body integrity. *Int J Med Sci* 2004;1(3):170-180.

Briggs AM, Greig AM, Wark JD. The vertebral fracture cascade in osteoporosis: a review of aetiopathogenesis. *Osteoporos Int* 2007 May;18(5):575-84. Review.

Duan Y, Seeman E, Turner CH. The biomechanical basis of vertebral body fragility in men and women. *J Bone Miner Res*. 2001 Dec;16(12):2276-83.

Edmondston SJ, Singer KP, Day RE, Price RI, Bredahl PD. Ex vivo estimation of thoracolumbar vertebral body compressive strength: the relative contributions of bone densitometry and vertebral morphometry. *Osteoporos Int* 1997;7(2):142-8.

Fonseca H, Moreira-Goncalves D, Coriolano HJ, Duarte JA. Bone quality: the determinants of bone strength and fragility. *Sports Med* 2014;44(1):37-53.

Fuchs RK, Kersh ME, Carballido-Gamio J, Thompson WR, Keyak JH, Warden SJ. Physical Activity for Strengthening Fracture Prone Regions of the Proximal Femur. *Curr Osteoporos Rep*. 2017;15(1):43-52. doi:10.1007/s11914-017-0343-6

Giangregorio, L. and R. El-Kotob (2017). "Exercise, muscle, and the applied load-bone strength balance." *Osteoporos Int* 28(1): 21-33.

Hayes WC, Piazza SJ, Zysset PK. Biomechanics of fracture risk prediction of the hip and spine by quantitative computed tomography. *Radiol Clin North Am* 1991Jan;29(1):1-18. Review.

Hsieh YF, Robling AG, Ambrosius WT, Burr DB, Turner CH. Mechanical loading of diaphyseal bone in vivo: the strain threshold for an osteogenic response varies with location. *J Bone Miner Res* 2001;16:2291-2297.

Hsieh YF, Turner CH. Effects of loading frequency on mechanically induced bone formation. *J Bone Miner Res* 2001;16:918-924.

Martelli S, Kersh ME, Schache AG, & Pandy MG. Strain energy in the femoral neck during exercise. *J Biomech* 2014;47(8), 1784-1791.

Myers ER, Wilson SE. Biomechanics of osteoporosis and vertebral fracture. *Spine* 1997 Dec 15;22(24 Suppl):25S-31S. Review.

Robling AG, Burr DB, Turner CH. Recovery periods restore mechanosensitivity to dynamically loaded bone. *J Exp Biol* 2001;204:3389-3399.

Robling AG, Hinant FM, Burr DB, Turner CH. Improved bone structure and strength after long-term mechanical loading is greatest if loading is separated into short bouts. *J Bone Miner Res*. 2002;17(8):1545-1554.

Robling AG, Duijvelaar KM, Geevers JV, Ohashi N, Turner CH. Modulation of appositional and longitudinal bone growth in the rat ulna by applied static and dynamic force. *Bone* 2001;29:105-113.

Rubin C, Turner AS, Muller R, et al. Quantity and quality of trabecular bone in the femur are enhanced by a strongly anabolic, noninvasive mechanical intervention. *J Bone Miner Res* 2002;17(2):349-357.

Saxon LK, Robling AG, Alam I, Turner CH. Mechanosensitivity of the rat skeleton decreases after a long period of loading, but is improved with time off. *Bone* 2005;36:454-464.

Schriefer JL, Warden SJ, Saxon LK, Robling AG, Turner CH. Cellular accommodation and the response of bone to mechanical loading. *J. Biomech* 2005;38:1838-1845.

Seeman E. Periosteal bone formation: a neglected determinant of bone strength. *N Engl J Med* 2003; 349:320-3.

Singer K, Edmondston S, Day R, Breidahl P, Price R. Prediction of thoracic and lumbar vertebral body compressive strength: correlations with bone mineral density and vertebral region. *Bone* Aug 1995;17(2): 167-74.

Turner CH. Toward a mathematical description of bone biology: the principal of cellular accommodation. *Calcif Tissue Int* 1999;65:466-471.

Warden SJ, Turner CH. Mechanotransduction in cortical bone is most efficient at loading frequencies of 5-10 Hz. *Bone* 2004;34:261-270.

Warden SJ, Hurst JA, Sanders MS, Turner CH, Burr DB, Li J. Bone adaptation to a mechanical loading program significantly increases skeletal fatigue resistance. *J Bone Miner Res* 2005 May;20(5):809-816.

Yang RS, Lin HJ, Chieng PU, Liu TK, Tsai KS. Estimated risk score for spine fracture in the specific bending activity of normal Taiwanese men and women. *Spine* 2005 Oct 15;30(20):2288-92.

### **3. IN VIVO MODELS RELATED TO FRACTURE RISK AND TO EXERCISE**

Andersson GB, Ortengren R, Nachemson A. Intradiskal pressure, intra-abdominal pressure and myoelectric back muscle activity related to posture and loading. *Clin Orthop Relat Res* 1977 NovDec;(129):156-64.

Briggs AM, Wrigley TV, van Dieën JH, Phillips B, Lo SK, Greig AM, Bennell KL. The effect of osteoporotic vertebral fracture on predicted spinal loads in vivo. *Eur Spine J* 2006 Dec;15(12):1785-95.

Sato K, Kikuchi S, Yonezawa T. In vivo intradiscal pressure measurement in healthy individuals and in patients with ongoing back problems. *Spine* 1999 Dec 1;24(23):2468-74.



Schultz AB. Analysis and measurement of lumbar trunk loads in tasks involving bends and twists. *J Biomech* 1982;15(9):669-75.

Torrance AG, Mosley JR, Suswillo RFL, Lanyon LE. Noninvasive loading of the rat ulna *in vivo* induces a strain-related modeling response uncomplicated by trauma or periosteal pressure. *Calcif Tissue Int* 1994;54:241-247.

Umemura Y, Ishiko T, Yamauchi T, Kurono M, Mashiko S. Five jumps per day increase bone mass and breaking force in rats. *J Bone Miner Res* 1997;12:1480-1485.

#### **4. PREDICTORS AND EFFECTS OF THORACIC HYPERKYPHOSIS**

Bartynski WS, Heller MT, Grahovac SZ, Rothfus WE, Kurs-Lasky M. Severe thoracic kyphosis in the older patient in the absence of vertebral fracture: Association of extreme curve with age. *Am J Neuroradiol.* 2005;26:2077-85.

Briggs AM, van Dieën JH, Wrigley TV, Greig AM, Phillips B, Lo SK, Bennell KL. Thoracic kyphosis affects spinal loads and trunk muscle force. *Phys Ther* 2007 May;87(5):595-607.

Bruno AG, Anderson DE, D'Agostino J, Boussein ML. The effect of thoracic kyphosis and sagittal plane alignment on vertebral compressive loading. *Journal of Bone and Mineral Research* 2012;27:2144-51.

Capponi D, Hlebs S. The impact of physical activity on the severity of spine curvatures and balance disorders in postmenopausal women. *Slovenian Nursing Review* 2008 06;42:77-86.

Culham EG, Jimenez HA, King CE. Thoracic kyphosis, rib mobility, and lung volumes in normal women and women with osteoporosis. *Spine* 1994 Jun 1;19(11):1250-5.

Di Bari M. Thoracic kyphosis and ventilatory dysfunction in unselected older persons: an epidemiological study in Dicomano, Italy. *J Am Geriatr Soc* 2004 Jun;52(6):909-15.

Eagan MS, Sedlock DA. Kyphosis in active and sedentary postmenopausal women. *Med Sci Sports Exerc* 2001 May;33(5):688-95.

Ensrud KE, Black DM, Harris F, Ettinger B, Cummings SR. Correlates of kyphosis in older women. The Fracture Intervention Trial Research Group. *J Am Geriatr Soc* 1997 Jun;45(6):682-7.

Ettinger B, Black DM, Palermo L, Nevitt MC, Melnikoff S, Cummings SR. Kyphosis in older women and its relation to back pain, disability and osteopenia: the study of osteoporotic fractures. *Osteoporos Int* 1994 Jan;4(1):55-60.

Fon GJ, Pitt MJ, Thies Jr. AC. Thoracic kyphosis: Range in normal subjects. *Am J Roentgenol* 1980;134:979-83.

Granito RN, Aveiro MC, Renno ACM, Oishi J, Driusso P. Comparison of thoracic kyphosis degree, trunk muscle strength and joint position sense among healthy and osteoporotic elderly women: A cross-sectional preliminary study. *Arch Gerontol Geriatr* 2012;54:e199-202.

Greendale GA, Nili NS, Huang MH, Seeger L, Karlamangla AS. The reliability and validity of three non-radiological measures of thoracic kyphosis and their relations to the standing radiological Cobb angle. *Osteoporosis Int* 2011;22:1897-905.

Greig AM, Briggs AM, Bennell KL, Hodges PW. Trunk muscle activity is modified in osteoporotic vertebral fracture and thoracic kyphosis with potential consequences for vertebral health. 2014; *PLoS One* 9(10): e109515.

Hinman MR. Comparison of thoracic kyphosis and postural stiffness in younger and older women. *Spine J* 2004 Jul-Aug;4(4):413-7.

Huang M, Barrett-Connor G, Greendale GA, Kado DM. Hyperkyphotic posture and risk of future osteoporotic fractures: The Rancho Bernardo study. *Journal of Bone and Mineral Research* 2006;21:419-23.

Kado DM, Huang M-, Karlamangla AS, Barrett-Connor E, Greendale GA. Hyperkyphotic posture predicts mortality in older community-dwelling men and women: A prospective study. *J Am Geriatr Soc* 2004;52:1662-7.

Kado DM, Huang MH, Barrett-Connor E, Greendale GA. Hyperkyphotic posture and poor physical functional ability in older community-dwelling men and women: the Rancho Bernardo study. *J Gerontol A Biol Sci Med Sci* May 2005;60(5):633-637.

Kado DM, Huang MH, Nguyen CB, Barrett-Connor E, Greendale GA. Hyperkyphotic posture and risk of injurious falls in older persons: the Rancho Bernardo Study. *J Gerontol A Biol Sci Med Sci* Jun 2007;62(6):652-657.

Kado DM, Miller-Martinez D, Lui LY, Cawthon P, Katzman WB, Hillier TA, Fink HA, Ensrud KE. Hyperkyphosis, Kyphosis Progression, and Risk of Non-Spine Fractures in Older Community Dwelling Women: The Study of Osteoporotic Fractures (SOF). *J Bone Miner Res* 2014 Oct; 29(10):2210-6.

Kado DM, Prenovost K, Crandall C. Narrative review: Hyperkyphosis in older persons. *Ann Intern Med* 2007;147:330-8.

Katzman WB, Huang MH, Lane NE, Ensrud KE, Kado DM. Kyphosis and decline in physical function over 15 years in older community-dwelling women: the Study of Osteoporotic Fractures. *J Gerontol A Biol Sci Med Sci*. 2013 Aug;68(8):976-83.

Katzman WB, Harrison SL, Fink HA, Marshall LM, Orwoll E, Barrett-Connor E, Cawthon PM, Kado DM; for the Osteoporotic Fractures in Men (MrOS) Study Research Group. Physical function in older men with hyperkyphosis. *J Gerontol A Biol Sci Med Sci*. 2014 Nov 27.

Katzman WB, Vittinghoff E, Kado DM, Lane NE, Ensrud KE, Shipp K. Thoracic kyphosis and rate of incident vertebral fractures: The Fracture Intervention Trial. *Osteoporosis Int*. 2016 Jan 18.

Katzman WB, Vittinghoff E, Kado DM, Schafer AL, Wong SS, Gladin A, Lane NE. Study of Hyperkyphosis, Exercise and Function (SHEAF) Protocol of a Randomized Controlled Trial of Multimodal Spine-Strengthening Exercise in Older Adults With Hyperkyphosis. *Phys Ther*. 2015 Aug 6.



Katzman WB, Vittinghoff E, Kado DM. Hyperkyphosis, Independent of Spinal Osteoporosis, is Associated with Impaired Mobility in Older Community-Dwelling Women. *Osteoporos Intl* 2010. May 18.

Katzman WB, Wanek L, Shepherd JA, Sellmeyer DE. Age-Related Hyperkyphosis: Its Causes and Consequences and Management. *J Orthop Sports Phys Ther* 2010 Jun;40(6):352-60. Review.

Katzman W, Cawthon P, Hicks GE, Vittinghoff E, Shepherd J, Cauley JA, et al. Association of spinal muscle composition and prevalence of hyperkyphosis in healthy community-dwelling older men and women. *J Gerontol A Biol Sci Med Sci* 2011;67A:191-195.

Keller TS, Harrison DE, Colloca CJ, Harrison DD, Janik TJ. Prediction of osteoporotic spinal deformity. *Spine* Mar 1 2003;28(5):455-46

Kusano M, et al. Size of hiatus hernia correlates with severity of kyphosis, not with obesity, in elderly Japanese women. *J Clin Gastroenterol* 2008 Apr;42(4):345-50.

Lewiecki EM. Kyphosis and vertebral fractures: What do we know now? *Aging Health*. 2009;5:483-7.

Lind LR, Lucente V, Kohn N. Thoracic kyphosis and the prevalence of advanced uterine prolapse. *Obstet Gynecol* 1996 Apr;87(4):605-9.

Mattox TF, et al. Abnormal spinal curvature and its relationship to pelvic organ prolapse. *Am J Obstet Gynecol* 2000 Dec;183(6):1381-4; discussion 1384.

Milne JS, Lauder IJ. Age effects in kyphosis and lordosis in adults. *Ann Hum Biol* 1974 Jul;1(3):327-37.

Nardo, L, Lane, NE, Parimi, N, Cawthon, P, Fan, B, Shepherd, J, Cauley, J, Zucker-Levine, A, Murphy, R, Katzman, WB. Diffuse Idiopathic Skeletal Hyperostosis Association with Thoracic Spine Kyphosis: A Cross-sectional Study for the Health Aging and Body Composition Study. *Spine*. 2014. Nov 15;39(24):E1418-24.

[Katzman WB, et al. Reliability of sagittal vertical axis measurement and association with measures of age-related hyperkyphosis. \*J Phys Ther Sci\*. 2018 Dec;30\(12\):1417-1423.](#)

Perriman DM, Scarvell JM, Hughes AR, Lueck CJ, Dear KBG, Smith PN. Thoracic hyperkyphosis: A survey of Australian physiotherapists. *Physiotherapy Research International* 2012;17:167-178.

Porter RW, Johnson K, McCutchan JD. Wrist fracture, heel bone density and thoracic kyphosis: a case control study. *Bone* 1990;11(3):211-4.

Puche RC, Morosano M, Masoni A, Jimeno NP, Bertoluzzo SM, Podadera JC, et al. The natural history of kyphosis in postmenopausal women. *Bone* 1995 17:239-46.

[Roghani T, Khalkhali Zavieh M, Rahimi A, Talebian S, Manshadi FD, Akbarzadeh Baghban A, Katzman W. \(2018\). The reliability and validity of a designed setup for the assessment of static back extensor force and endurance in older women with and without hyperkyphosis. \*Physiother Theory Pract\*, 34\(11\), 882-893.](#)

Ryan SD, Fried LP. The impact of kyphosis on daily functioning. *J Am Geriatr Soc* 1997;45:1479-86.

Schneider DL, Von Mühlen DG, Barrett-Connor E, Sartoris DJ. Kyphosis does not equal vertebral fractures: The Rancho Bernardo study. *J Rheumatol* 2004;31:747-52.

Schuerman S. Relationships between postural exercise and risk factors for falling in individuals with osteoporosis [dissertation]. Omaha: University of Nebraska;1998.

Sinaki M, Brey RH, Hughes CA, Larson DR, Kaufman KR. Balance disorder and increased risk of falls in osteoporosis and kyphosis: Significance of kyphotic posture and muscle strength. *Osteoporosis Int* 2005;16:1004-10.

Takahashi T, Ishida K, Hirose D, Nagano Y, Okumiya K, Nishinaga M, et al. Trunk deformity is associated with a reduction in outdoor activities of daily living and life satisfaction in community-dwelling older people. *Osteoporosis Int* 2005;16:273-9.

Teramoto S, Suzuki M, Matsuse T, Ohga E, Katayama H, Nagase T, et al. Influence of kyphosis on the age-related decline in pulmonary function. *Japanese Journal of Geriatrics* 380 1998;35:23-7.

Vialle R, et al. Radiographic analysis of the sagittal alignment and balance of the spine in asymptomatic subjects. *Bone Joint Surg Am* 2005 Feb;87(2):260-7.

Voutsinas SA, MacEwen GD. Sagittal profiles of the spine. *Clin Orthop* 1986:235-42.

## **5. EXERCISE INTERVENTIONS FOR KYPHOSIS AND/OR VERTEBRAL FRACTURE**

Ball JM, Cagle P, Johnson BE, Lucasey C, Lukert BP. Spinal extension exercises prevent natural progression of kyphosis. *Osteoporosis Int* 2009 Mar;20(3):481-9.

Bansal S, Katzman WB, Giangregorio LM. Exercise for improving age-related hyperkyphotic posture: a systematic review. *Arch Phys Med Rehabil* 2014 Jan; 95(1):129-40.

Barker, KL. Exercise or manual physiotherapy compared with a single session of physiotherapy for osteoporotic vertebral fracture: three-arm PROVE RCT. *Health Technol Assess*, 2019;23(44)

Barker, KL, Javaid, MK, Newman M, Minns Lowe C, Stallard N, Campbell H, Lamb S. Physiotherapy Rehabilitation for Osteoporotic Vertebral Fracture (PROVE): study protocol for a randomized controlled trial. *Trials* 2014;15, 22.

Barker KL, Newman M, Stallard N, Leal J, Lowe CM, Javaid MK, Noufaily A, Hughes T, Smith D, Gandhi V, Cooper C, Lamb SE; PROVE trial group. Physiotherapy rehabilitation for osteoporotic vertebral fracture-a randomised controlled trial and economic evaluation (PROVE trial). *Osteoporosis Int* 2020 Feb;31(2):277-289. <http://doi.org/10.1007/s00198-019-05133-0>.

Bautmans I, Van Arken J, Van Mackelenberg M, Mets T. Rehabilitation using manual mobilization for thoracic kyphosis in elderly postmenopausal patients with osteoporosis. *J Rehabil Med* 2010;42:129-35.

Bello M, Sousa, MC, Neto G, Oliveira L, Guerras I, Mendes R, Sousa N. The effect of a long-term, community-based exercise program on bone mineral density in postmenopausal women with pre-diabetes and type 2 diabetes. *J Hum Kinet* 2014; 43: 43-48.

Benedetti MG, Berti L, Presti C, Frizziero A, Giannini S. Effects of an adapted physical activity program in a group of elderly subjects with flexed posture: Clinical and instrumental assessment. *Journal of NeuroEngineering and Rehabilitation* 2008;5:32.

Bennell KL, Matthews B, Greig A, Briggs A, Kelly A, Sherburn M, Larsen J, Wark J. Effects of an exercise and manual therapy program on physical impairments, function and quality-of-life in people with osteoporotic vertebral fracture: a randomised, single-blind controlled pilot trial. *BMC Musculoskelet Disord* 2010 Feb 17;11:36.

Bergland A, Thorsen H, Karesen R. Effect of exercise on mobility, balance, and health-related quality of life in osteoporotic women with a history of vertebral fracture: a randomized, controlled trial. *Osteoporos Int* 2011;22:1863-1871.

Bergstrom I, Bergstrom K, Kronhed A-C, et al. Back extensor training increases muscle strength in postmenopausal women with osteoporosis, kyphosis, and vertebral fractures. *Advances in Physiotherapy* 2011;13:110-117.

Evstigneeva L, Lesnyak O, Bultink IE, et al. Effect of twelve-month physical exercise program on patients with osteoporotic vertebral fractures: a randomized, controlled trial. *Osteoporos Int*. 2016;27(8):2515-2524. doi:10.1007/s00198-016-3560-4

Giangregorio LM, Papaioannou A, Macintyre NJ, et al. Too Fit To Fracture: exercise recommendations for individuals with osteoporosis or osteoporotic vertebral fracture. *Osteoporos Int* 2014;25(3):821-835.

[Giangregorio LM](#)<sup>1</sup>, [McGill S](#), [Wark JD](#), [Laprade J](#), [Heinonen A](#), [Ashe MC](#), [MacIntyre NJ](#), [Cheung AM](#), [Shipp K](#), [Keller H](#), [Jain R](#), [Papaioannou A](#). Too Fit To Fracture: outcomes of a Delphi consensus process on physical activity and exercise recommendations for adults with osteoporosis with or without vertebral fractures. *Osteoporos Int* 2015; Mar;26(3):891-910.

[Giangregorio LM](#), et al. Build better bones with exercise (B3E pilot trial): results of a feasibility study of a multicenter randomized controlled trial of 12 months of home exercise in older women with vertebral fracture. *Osteoporos Int*, 2018;29(11), 2545-2556.

Gold DT, Shipp KM, Pieper CF, et al. Group treatment improves trunk strength and psychological status in older women with vertebral fractures: results of a randomized, clinical trial. *J Am Geriatr Soc* 2004;54:1471-1478.

Greendale GA, Huang M-, Karlamangla AS, Seeger L, Crawford S. Yoga decreases kyphosis in senior women and men with adult-onset hyperkyphosis: Results of a randomized controlled trial. *J Am Geriatr Soc* 2009;57:1569-79.

Greendale GA, McDivit A, Carpenter A, Seeger L, Huang M. Yoga for women with hyperkyphosis: Results of a pilot study. *Am J Public Health* 2002;92:1611-4.

Greig AM, Bennell KL, Briggs AM, Hodges PW. Postural taping decreases thoracic kyphosis but does not influence trunk muscle EMG activity or balance in women with osteoporosis. *Man Ther* 2008;13:249-57.

Hakestad KA, Torstveit, M. K, Nordsletten L, Axelsson AC, Risberg MA. Exercises Including Weight Vests and a Patient Education Program for Women With Osteopenia: A Feasibility Study of the OsteoACTIVE Rehabilitation Program. *J Orthop Sports Phys Ther* 2015; 45(2): 97-105.

Huntoon EA, Schmidt CK, Sinaki M. Significantly fewer refractures after vertebroplasty in patients who engage in back-extensor-strengthening exercises. *Mayo Clin Proc* 2008 Jan;83(1):54-7.

Itoi E, Sinaki M. Effect of back-strengthening exercise on posture in healthy women 49 to 65 years of age. *Mayo Clin Proc* 1994 Nov;69(11):1054-9.

Jang HJ, Hughes LC, Oh DW, & Kim SY. Effects of Corrective Exercise for Thoracic Hyperkyphosis on Posture, Balance, and Well-Being in Older Women: A Double-Blind, Group-Matched Design. *J Geriatr Phys Ther* 2019;42(3), E17-e27.

Kado DM. The rehabilitation of hyperkyphotic posture in the elderly. *European Journal of Physical and Rehabilitation Medicine* 2009;45:583-93.

Katzman WB, Gladin A, Lane NE, et al. Feasibility and Acceptability of Technology-Based Exercise and Posture Training in Older Adults With Age-Related Hyperkyphosis: Pre-Post Study. *JMIR Aging*. 2019;2(1):e12199. doi:10.2196/12199

Katzman WB, Sellmeyer DE, Stewart AL, et al. Changes in flexed posture, musculoskeletal impairment, and physical performance after group exercise in community-dwelling older women. *Arch Phys Med Rehabil* 2007;88:192-199.

Lu YH, et al. (2016). "Twelve-Minute Daily Yoga Regimen Reverses Osteoporotic Bone Loss." Topics in Geriatric Rehabilitation, *Top Geriatr Rehabil*. 2016;32(2):81–87.

Marini S, Leoni E, Raggi A, et al. Proposal of an Adapted Physical Activity Exercise Protocol for Women with Osteoporosis-Related Vertebral Fractures: A Pilot Study to Evaluate Feasibility, Safety, and Effectiveness. *Int J Environ Res Public Health*. 2019;16(14):2562. Published 2019 Jul 18. doi:10.3390/ijerph16142562

Olsen CF & Bergland A. The effect of exercise and education on fear of falling in elderly women with osteoporosis and a history of vertebral fracture: results of a randomized controlled trial. *Osteoporos Int* 2014;25(8), 2017-2025.

Papaioannou A, Adachi JD, Winegard K et al. Efficacy of home-based exercise for improving quality of life among elderly women with symptomatic osteoporosis-related vertebral fracture *Osteoporos Int* 2003;14:677-682.

Pawlowsky SB, Hamel KA, Katzman WB. Stability of kyphosis, strength, and physical performance gains 1 year after a group exercise program in community-dwelling hyperkyphotic older women. *Arch Phys Med Rehabil* 2009 Feb;90(2):358-61.

Renno ACM, Granito RN, Driusso P, Costa D, Oishi J. Effects of an exercise program on respiratory function, posture and on quality of life in osteoporotic women: A pilot study. *Physiotherapy* 2005;91:113-8.

Shipp KM, Gold DT, Pieper CF, Lyles KW. Improving Trunk Strength and Endurance in Older Women with Vertebral Fractures. *J Bone Miner Res* 2010;22:S463.

Sinaki M, Wollan PC, Scott RW, Gelczer RK. Can strong back extensors prevent vertebral fractures in women with osteoporosis? *Mayo Clin Proc* 1996 Oct;71(10):951-6.

Sinaki M, Itoi E, Wahner HW, et al. Stronger back muscles reduce the incidence of vertebral fractures: a prospective 10 year follow-up of postmenopausal women. *Bone* 2002;30(6):836-841.

Sinaki M, Brey RH, Hughes CA, Larson DR, Kaufman KR. Significant reduction in risk of falls and back pain in osteoporotic-kyphotic women through a Spinal Proprioceptive Extension Exercise Dynamic (SPEED) program. *Mayo Clin Proc* 2005 Jul;80(7):849-55.

Wang R-, Wang Y, Bai J-, Guo J, Wang J-, Chai W, et al. Effects of rehabilitation exercise on pulmonary function, thoracic kyphosis and physical activity in old female patients with osteoporosis. *Chinese Journal of Clinical Rehabilitation*.2006;10:34-6.

Yang I, He CQ, Xie W, et al. Effect of pain-free exercises on female osteoporosis patients with spinal compression fracture. *J Clin Rehabil Tissue Eng Res* 2007;11:9108-9111.

## **6.EPIDEMIOLOGY AND RISK OF FRACTURE**

Alghadir AH, Aly FA, & Gabr SA. Effect of Moderate Aerobic Training on Bone Metabolism Indices among Adult Humans. *Pak J Med Sci* 2014;30(4), 840-844.

Bartalena T, Rinaldi MF, Modolon C, et al. Incidental vertebral compression fractures in imaging studies: Lessons not learned by radiologists. *World J Radiol.* 2010;2(10):399-404.  
doi:10.4329/wjr.v2.i10.399

Boudreau DM, Yu O, Balasubramanian A, et al. A Survey of Women's Awareness of and Reasons for Lack of Postfracture Osteoporotic Care. *J Am Geriatr Soc.* 2017;65(8):1829-1835.  
doi:10.1111/jgs.14921

Caban-Martinez AJ, Courtney TK, Chang WR, et al. Leisure-Time Physical Activity, Falls, and Fall Injuries in Middle-Aged Adults. *Am J Prev Med.* 2015;49(6):888-901.  
doi:10.1016/j.amepre.2015.05.022

Chahal J, Lee R, & Luo J. Loading dose of physical activity is related to muscle strength and bone density in middle-aged women. *Bone* 2014;67C, 41-45.

Dallas SL, Prideaux M, Bonewald LF. The osteocyte: an endocrine cell ... and more. *Endocr Rev* 2013; Oct;34(5):658-90.

Delmas PD, Genant HK, Crans GG, et al. Severity of prevalent vertebral fractures and the risk of subsequent vertebral and nonvertebral fractures: results from the MORE trial. *Bone* 2003;33(4):522-532.

Eisman, JA, Bogoch ER, Dell R, Harrington JT, McKinnet RE, McLellan A, Mitchell PJm Silverman S, Singleton R, Siris E ASBMR Task Force On Secondary Fracture Prevention: Making the first fracture the last: ASBMR task force report on secondary fracture prevention. *J Bone Mineral Res* 2012; Oct 27 (10):2039-46

Evans R, Negus C, Centi A et al. Peripheral QCT Sector Analysis Reveals Exercise-induced Increases in Tibial Bone Mineral Density. *J Musculoskelet Neuronal Interact* 2012;12:155-64.

[Huang CY, Liao LC, Tong KM, et al. Mediating effects on health-related quality of life in adults with osteoporosis: a structural equation modeling. \*Osteoporos Int.\* 2015;26\(3\):875-883. doi:10.1007/s00198-014-2963-3](#)

Kanis JA, Johnell O, De Laet C, Johansson H, Oden A, Delmas P, Eisman J, Fujiwara S, Garnero P, Kroger H, McCloskey EV, Mellstrom D, Melton LJ, Pols H, Reeve J, Silman A, Tenenhouse A. A meta-analysis of previous fracture and subsequent fracture risk. *Bone* 2004 Aug;35(2):375-82.

Kontulainen S, Sievanen H, Kannus P, Pasanen M, Vuori I. Effect of long-term impact-loading on mass, size, and estimated strength of humerus and radius of female racquet-sports players: a peripheral quantitative CT study between young and old starters and controls. *J Bone Miner Res* 2003;18:352-9.

Leech JA, Dulberg C, Kellie S, Pattee L, Gay J. Relationship of lung function to severity of osteoporosis in women. *Am Rev Respir Dis* 1990 Jan;141(1):68-71.

[Li Y, Yan L, Cai S, Wang P, Zhuang H, Yu H. The prevalence and under-diagnosis of vertebral fractures on chest radiograph. \*BMC Musculoskelet Disord.\* 2018;19\(1\):235. Published 2018 Jul 18. doi:10.1186/s12891-018-2171-y](#)

Lindsay R, Silverman SL, Cooper C, et al. Risk of new vertebral fracture in the year following a fracture. *JAMA.* 2001;285(3):320-323.

Morseth B, Ahmed L, Djornerem A et al. Leisure Time Physical Activity and Risk of Non-vertebral Fracture in Men and Women Age 55 Years and Older: The Tromso Study. *Euro J Epidemiol* 2012.

[Raybould G, Babatunde O, Evans AL, Jordan JL, Paskins Z. Expressed information needs of patients with osteoporosis and/or fragility fractures: a systematic review. \*Arch Osteoporos.\* 2018;13\(1\):55. Published 2018 May 8. doi:10.1007/s11657-018-0470-4](#)

Sawka AM, Thabane L, Papaioannou A, et al. Health-related quality of life measurements in elderly Canadians with osteoporosis compared to other chronic medical conditions: a population-based study from the Canadian Multicentre Osteoporosis Study (CaMos). *Osteoporos Int.* 2005;16(12):1836-1840. doi:10.1007/s00198-005-1949-6



Singer A, Exuzides A, Spangler L, et al. Burden of illness for osteoporotic fractures compared with other serious diseases among postmenopausal women in the United States. *Mayo Clin Proc.* 2015;90(1):53-62. doi:10.1016/j.mayocp.2014.09.011

van Geel TA, van Helden S, Geusens PP, Winkens B, Dinant GJ. Clinical subsequent fractures cluster in time after first fractures. *Ann Rheum Dis.* 2009;68(1):99-102. doi:10.1136/ard.2008.092775

Vokó Z, Gáspár K, Inotai A, et al. Osteoporotic fractures may impair life as much as the complications of diabetes. *J Eval Clin Pract.* 2017;23(6):1375-1380. doi:10.1111/jep.12800

Wang M, Wu F, Callisaya ML, Jones G, Winzenberg T. Incidence and circumstances of falls among middle-aged women: a cohort study [published online ahead of print, 2020 Sep 12]. *Osteoporos Int.* 2020;10.1007/s00198-020-05617-4. doi:10.1007/s00198-020-05617-4

Wasnich RD, Davis JW, Ross PD. Spine fracture risk is predicted by non-spine fractures. *Osteoporos Int* 1994 Jan;4(1):1-5.

Weaver J, Sajjan S, Lewiecki EM, Harris ST, Marvos P. Prevalence and Cost of Subsequent Fractures Among U.S. Patients with an Incident Fracture. *J Manag Care Spec Pharm.* 2017;23(4):461-471. doi:10.18553/jmcp.2017.23.4.461

## **7. FLEXION EXERCISES**

Sinaki M, Mikkelsen BA. Postmenopausal spinal osteoporosis: flexion versus extension exercises. *Arch Phys Med Rehabil* 1984 Oct;65(10):593-6.

Sinaki M. Yoga spinal flexion positions and vertebral compression fracture in osteopenia or osteoporosis of spine: case series. *Pain Pract* 2013 Jan;13(1):68-75.

## **8. OTHER EXERCISE AND ACTIVITY INTERVENTIONS, ASSESSMENTS, & MEDICAL MANAGEMENT FOR BONE HEALTH**

Aboarrage Junior, AM, et al. A High-Intensity Jump-Based Aquatic Exercise Program Improves Bone Mineral Density and Functional Fitness in Postmenopausal Women. *Rejuvenation Res* 2018;21(6):535-540.

Agostini D, Zeppa Donati S, Lucertini F, et al. Muscle and Bone Health in Postmenopausal Women: Role of Protein and Vitamin D Supplementation Combined with Exercise Training. *Nutrients.* 2018;10(8):1103. Published 2018 Aug 16. doi:10.3390/nu10081103

Anastasilakis AD, Polyzos SA, Makras P, Aubry-Rozier B, Kaouri S, Lamy O. Clinical Features of 24 Patients With Rebound-Associated Vertebral Fractures After Denosumab Discontinuation: Systematic Review and Additional Cases. *J Bone Miner Res.* 2017;32(6):1291-1296. doi:10.1002/jbmr.3110

Bolton KL, Egerton T, Wark J, et al. Effects of exercise on bone density and falls risk factors in post-menopausal women with osteopenia: a randomised controlled trial. *J Sci Med Sport* 2012;15(2):102-109.

Camacho PM, Petak SM, Binkley N, et al. AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS/AMERICAN COLLEGE OF ENDOCRINOLOGY CLINICAL PRACTICE GUIDELINES FOR THE DIAGNOSIS AND TREATMENT OF POSTMENOPAUSAL OSTEOPOROSIS-2020 UPDATE. *Endocr Pract.* 2020;26(Suppl 1):1-46. doi:10.4158/GL-2020-0524SUPPL

Conley RB, Adib G, Adler RA, et al. Secondary Fracture Prevention: Consensus Clinical Recommendations from a Multistakeholder Coalition. *J Bone Miner Res.* 2020;35(1):36-52. doi:10.1002/jbmr.3877

Compston J, Cooper A, Cooper C, et al. UK clinical guideline for the prevention and treatment of osteoporosis. *Arch Osteoporos.* 2017;12(1):43. doi:10.1007/s11657-017-0324-5

Cummings SR, Ferrari S, Eastell R, et al. Vertebral Fractures After Discontinuation of Denosumab: A Post Hoc Analysis of the Randomized Placebo-Controlled FREEDOM Trial and Its Extension. *J Bone Miner Res.* 2018;33(2):190-198. doi:10.1002/jbmr.3337

Curtis J, Silverman S, The Five Ws of a Fracture Liaison Service: Why, Who, What, Where and How? in Osteoporosis, We Reap What We Sow; *Curr Osteoporosis Rep* 2013;11:365-368.

Ducher G, Bass SL, Saxon L, Daly RM. Effects of repetitive loading on the growth-induced changes in bone mass and cortical bone geometry: a 12-month study in pre/peri-and postmenarcheal tennis players. *J Bone Miner Res* 2011; 26:1321-9.

Ebeling PR, Akesson K, Bauer DC, et al. The Efficacy and Safety of Vertebral Augmentation: A Second ASBMR Task Force Report. *J Bone Miner Res.* 2019;34(1):3-21. doi:10.1002/jbmr.3653

Eisman JA, Bogoch ER, Dell R, et al. Making the first fracture the last fracture: ASBMR task force report on secondary fracture prevention. *J Bone Miner Res.* 2012;27(10):2039-2046. doi:10.1002/jbmr.1698

Gallacher SJ, Setting up an osteoporosis fracture liaison service: background and potential outcomes. *Best Practice Res Clin Rheumatol* 2005 Dec;19 (6):1081-94.

Garcia-Gomariz C, et al. Effect of 2 years of endurance and high-impact training on preventing osteoporosis in postmenopausal women: randomized clinical trial. *Menopause* 2018;25(3):301-306.

Giangregorio LM, MacIntyre NJ, Heinonen A, et al. Too Fit To Fracture: a consensus on future research priorities in osteoporosis and exercise. *Osteoporos Int* 2014;25(5):1465-1472.

Giangregorio, L M, et al. Build better bones with exercise: protocol for a feasibility study of a multicenter randomized controlled trial of 12 months of home exercise in women with a vertebral fracture. *Phys Ther* 2014;94(9):1337-1352.

Gianoudis J, Bailey CA, Sanders KM, et al. Osteo-cise: strong bones for life: protocol for a community-based randomised controlled trial of a multi-modal exercise and osteoporosis education program for older adults at risk of falls and fractures. *BMC Musculoskelet Disord* 2012;13:78.

Gianoudis J, Bailey CA, Ebeling PR, et al. Effects of a targeted multimodal exercise program incorporating high-speed power training on falls and fracture risk factors in older adults: a community-based randomized controlled trial. *J Bone Miner Res* 2014;29(1):182-191.

Giles M, Van Der Kallen J, Parker V, Cooper K, Ross L, McNeill SA. Team Approach: implementing a model of care for preventing osteoporosis related fractures; *Osteoporosis Int* 2011;22:2321-2328.

Gonzalo-Encabo P, et al. Dose-response effects of exercise on bone mineral density and content in post-menopausal women. *Scand J Med Sci Sports* 2019;29(8):1121-1129.

Gray M, Di Brezzo R, Fort IL. The effects of power and strength training on bone mineral density in premenopausal women. *J Sports Med Phys Fitness* 2013;53(4):428-436.

Greenway KG, Walkley JW, Rich PA. Does long-term swimming participation have a deleterious effect on the adult female skeleton? *Eur J Appl Physiol* 2012;112(9):3217-3225.

Harding AT, Weeks BK, Lambert C, Watson SL, Weis LJ, Beck BR. Effects of supervised high-intensity resistance and impact training or machine-based isometric training on regional bone geometry and strength in middle-aged and older men with low bone mass: The LIFTMOR-M semi-randomised controlled trial. *Bone*. 2020;136:115362. doi:10.1016/j.bone.2020.115362

Harding AT, Weeks BK, Lambert C, Watson SL, Weis LJ, Beck BR. A Comparison of Bone-Targeted Exercise Strategies to Reduce Fracture Risk in Middle-Aged and Older Men with Osteopenia and Osteoporosis: LIFTMOR-M Semi-Randomized Controlled Trial. *J Bone Miner Res*. 2020;35(8):1404-1414. doi:10.1002/jbmr.4008

Howe TE, Shea B, Dawson LJ, et al. Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Database of Systematic Reviews* 2011;issue 7. Art. No.: CD000333.

Kemmler W, Kohl M, Fröhlich M, et al. Effects of High-Intensity Resistance Training on Osteopenia and Sarcopenia Parameters in Older Men with Osteosarcopenia-One-Year Results of the Randomized Controlled Franconian Osteopenia and Sarcopenia Trial (FrOST). *J Bone Miner Res*. 2020;35(9):1634-1644. doi:10.1002/jbmr.4027

Kemmler K, von Stengel S, Bebnik M et al. Exercise and fractures in postmenopausal women: 12-year results of the Erlangen Fitness & Osteoporosis Prevention Study (EFOPS). *Osteopor Int* 2012;23:1267-76.

Kemmler W, Bebnik M, Kohl M, von Stengel S. "Exercise and fractures in postmenopausal women. Final (16 year) results of the controlled Erlangen Fitness and Osteoporosis Prevention Study (EFOPS)." *Osteoporos Int* 2015;12 May.

Kemmler W, & von Stengel S. Dose-response effect of exercise frequency on bone mineral density in post-menopausal, osteopenic women. *Scand J Med Sci Sports* 2014;24(3), 526-534.

Kerr C, Bottomley C, Shingler S, et al. The importance of physical function to people with osteoporosis. *Osteoporos Int*. 2017;28(5):1597-1607. doi:10.1007/s00198-017-3911-9

Lems WF, Dreinhöfer KE, Bischoff-Ferrari H, et al. EULAR/EFORT recommendations for management of patients older than 50 years with a fragility fracture and prevention of subsequent fractures. *Ann Rheum Dis*. 2017;76(5):802-810. doi:10.1136/annrheumdis-2016-210289

Lewiecki EM. Role of sclerostin in bone and cartilage and its potential as a therapeutic target in bone diseases. *Ther Adv Musculoskelet Dis* 2014;6(2), 48-57.

Lindsey C, Bookstein N. Physical Therapy for Osteoporosis: Prevention and Management. *American Physical Therapy Association: Section on Women's Health Home Study Course* 2013.

Majumdar SR, Kim N, Colman I, et al. Incidental vertebral fractures discovered with chest radiography in the emergency department: prevalence, recognition, and osteoporosis management in a cohort of elderly patients. *Arch Intern Med*. 2005;165(8):905-909. doi:10.1001/archinte.165.8.905

McArthur, C., et al. (2016). "Suggestions for Adapting Yoga to the Needs of Older Adults with Osteoporosis." *J Altern Complement Med* 22(3): 223-226.

Malmros B, Mortensen L, Jensen MB, Charles P. Positive effects of physiotherapy on chronic pain and performance in osteoporosis. *Osteoporos Int* 1998;8:215-221.

McNamara A, Gunter K, The influence of participation in Better Bones and Balance on skeletal health: evaluation of a community based program to reduce fall and fracture risk. *Osteoporosis Int* 2012; Jun:23 (6)1813-22.

McNamara AJ, Pavol MJ, Gunter KB. Meeting physical activity guidelines through community-based group exercise: "better bones and balance". *J Aging Phys Act* 2013;21(2):155-166.

Metcalf L, Lohman TG, Goings S, Houtkooper L, Ferreria D, Flint-Wagner H, Guido T, Martin J, Wright J, Cussler E. "Postmenopausal Women and Exercise for Prevention of Osteoporosis (BEST Study)." *ACSM's Health and Fitness Journal* 2001 5(3): 6-14.

Mitchell RM, Jewell P, Javaid MK, McKean D, Ostlere SJ. Reporting of vertebral fragility fractures: can radiologists help reduce the number of hip fractures?. *Arch Osteoporos*. 2017;12(1):71. doi:10.1007/s11657-017-0363-y

Miko I, et al. Effect of a balance-training programme on postural balance, aerobic capacity and frequency of falls in women with osteoporosis: A randomized controlled trial. *J Rehabil Med*, 2018;50(6), 542-547.

Multanen J, Nieminen, MT, Hakkinen A, Kujala UM, Jamsa T, Kautiainen H, Lammentausta E, Ahola R, Selanne H, Ojala R, Kiviranta I, Heinonen A. Effects of high-impact training on bone and articular cartilage: 12-month randomized controlled quantitative MRI study. *J Bone Miner Res* 2014;29(1), 192-201.

Murtezani A, Nevzati A, Ibraimi Z, Sllamniku S, Meka VS, Abazi N. The Effect of Land versus Aquatic Exercise Program on Bone Mineral Density and Physical Function in Postmenopausal Women with Osteoporosis: a Randomized Controlled Trial. *Ortop Traumatol Rehabil* 2014;16(3):319-325.

Newman M, Newman R, Hughes T, Vadher K, Barker KL. Is the timed loaded standing test a valid measure of back muscle endurance in people with vertebral osteoporosis?. *Osteoporos Int*. 2018;29(4):893-905. doi:10.1007/s00198-017-4358-8

Olenginski TP, Maloney-Saxon G, Matzko CK, et al. High-risk osteoporosis clinic (HiROC): improving osteoporosis and postfracture care with an organized, programmatic approach. *Osteoporos Int*. 2015;26(2):801-810. doi:10.1007/s00198-014-2967-z

Pasqualini L, Ministrini S, Lombardini R, et al. Effects of a 3-month weight-bearing and resistance exercise training on circulating osteogenic cells and bone formation markers in postmenopausal women with low bone mass. *Osteoporos Int*. 2019;30(4):797-806. doi:10.1007/s00198-019-04908-9

Paolucci T, Morone G, Iosa M, Grasso MR, Buzi E, Zangrando F, Paolucci S, Saraceni VM, Fusco A. Efficacy of group adapted physical exercises in reducing back pain in women with postmenopausal osteoporosis. *Aging Clin Exp Res* 2014;26(4), 395-402.

Pfeifer M, Kohlwey L, Begerow B, Minne HW. Effects of two newly developed spinal orthoses on trunk muscle strength, posture, and quality-of-life in women with postmenopausal osteoporosis: A randomized trial. *American Journal of Physical Medicine and Rehabilitation* 2011;90:805-15.

Qaseem A, Forcica MA, McLean RM, Denberg TD; Clinical Guidelines Committee of the American College of Physicians. Treatment of Low Bone Density or Osteoporosis to Prevent Fractures in Men and Women: A Clinical Practice Guideline Update From the American College of Physicians [published correction appears in *Ann Intern Med*. 2017 Sep 19;167(6):448]. *Ann Intern Med*. 2017;166(11):818-839. doi:10.7326/M15-1361

Rodriguez AJ, Fink HA, Mirigian L, et al. Pain, Quality of Life, and Safety Outcomes of Kyphoplasty for Vertebral Compression Fractures: Report of a Task Force of the American Society for Bone and Mineral Research. *J Bone Miner Res*. 2017;32(9):1935-1944. doi:10.1002/jbmr.3170

Saravi F, Sayegh F. Bone Mineral Density in Body Composition of Adult Premenopausal Women with Three Levels of Physical Activity. *J Osteoporos* 2013;953271.

Scottish Intercollegiate Guidelines Network (SIGN). Management of osteoporosis and the prevention of fragility fractures. Edinburgh: SIGN; 2015. (SIGN publication no. 142). [March 2015]. Available from URL: <http://www.sign.ac.uk>

Shipp, KM; Purse, JL; Gold, DT; Pieper, CF; Sloane, R; Schenkman, M; Lyles, KW. Timed loaded standing: a measure of combined trunk and arm endurance suitable for people with vertebral osteoporosis. *Osteoporosis International* 2002;11:914-922.

Speerin R, Slater H, Li L, Moore K, Chan M, Dreinhofer K, Ebeling PR, Willcock

S, Briggs, AM. Moving from evidence to practice: Models of care for the prevention and management of musculoskeletal conditions. *Best Pract Res Clin Rheumatol* 2014;28(3): 479-515.

Troy KL, Mancuso ME, Butler TA, Johnson JE. Exercise Early and Often: Effects of Physical Activity and Exercise on Women's Bone Health. *Int J Environ Res Public Health*. 2018;15(5):878. Published 2018 Apr 28. doi:10.3390/ijerph15050878

Troy KL, Mancuso ME, Johnson JE, Wu Z, Schnitzer TJ, Butler TA. Bone Adaptation in Adult Women Is Related to Loading Dose: A 12-Month Randomized Controlled Trial. *J Bone Miner Res*. 2020;35(7):1300-1312. doi:10.1002/jbmr.3999

Tu KN, Lie JD, Wan CKV, et al. Osteoporosis: A Review of Treatment Options. *P T*. 2018;43(2):92-104.

Ubago-Guisado E, et al. Effects of Zumba® and Aquagym on Bone Mass in Inactive Middle-Aged Women. *Medicina (Kaunas)*, 2019;55(1).

Watson SL, et al. Heavy resistance training is safe and improves bone, function, and stature in postmenopausal women with low to very low bone mass: novel early findings from the LIFTMOR trial. *Osteoporos Int*. 2015;26(12):2889-94.

Watson SL, et al. High-Intensity Resistance and Impact Training Improves Bone Mineral Density and Physical Function in Postmenopausal Women With Osteopenia and Osteoporosis: The LIFTMOR Randomized Controlled Trial. *J Bone Miner Res*. 2018;33(2):211-220.

Watson, SL, et al. High-intensity exercise did not cause vertebral fractures and improves thoracic kyphosis in postmenopausal women with low to very low bone mass: The LIFTMOR trial. *Osteoporos Int*. 2019;30(5):957-964.

Wu CH, Tu ST, Chang YF, et al. Fracture liaison services improve outcomes of patients with osteoporosis-related fractures: A systematic literature review and meta-analysis. *Bone*. 2018;111:92-100. doi:10.1016/j.bone.2018.03.018

## **9. HIP FRACTURE**

Binder EF, Brown M, Sinacore DR, et al. Effects of extended outpatient rehabilitation after hip fracture: a randomized controlled trial. *JAMA* 2004;292:837-846.

Bischoff-Ferrari HA, Dawson-Hughes B, Platz A, et al. Effect of high-dosage cholecalciferol and extended physiotherapy on complications after hip fracture: a randomized controlled trial. *Arch Intern Med* 2010;170:813-820.

Feskanich D, Willett W, Colditz G . Walking and leisure-time activity and risk of hip fracture in postmenopausal women. *JAMA* 2002;288(18):2300-2306.

Handoll HHG, Sherrington C, Mak JCS. Interventions for improving mobility after hip fracture surgery in adults. *Cochrane Database of Systematic Reviews* 2011;issue 3. Art No: CD001704.

Hauer K, Specht N, Schuler M, et al. Intensive physical training in geriatric patients after severe falls and hip surgery. *Age Aging* 2002;31:49-57.



Mangione KK, Craik RI, Palombaro KM, et al. Home-based leg-strengthening exercise improves function 1 year after hip fracture: a randomized controlled study. *J Am Geriatr Soc* 2010;58:1911-1917.

Mangione KK, Craik RI, Tomlinson SS, et al. Can elderly patients who have had a hip fracture perform moderate- to high-intensity exercise at home? *Phys Ther* 2005;85:727-739.

Orwig DI, Hochberg M, Yu-Yahiro J, et al. Delivery and outcomes of a yearlong home exercise program after hip fracture: a randomized controlled trial. *Arch Intern Med* 2011;171:323-331.

Portegijs E, Kallinen M, Rantanen T, et al. Effects of resistance training on lower-extremity impairments in older people with hip fracture. *Arch Phys Med Rehabil* 2008;89:1667-1674.

Sherrington C, Lord SR, Herbert RD. A randomized controlled trial of weight-bearing versus non-weight-bearing exercise for improving physical ability after usual care for hip fracture. *Arch Phys Med Rehabil* 2004;85:710-716.

Sherrington C, Lord SR. Home exercise to improve strength and walking velocity after hip fracture: a randomized controlled trial. *Arch Phys Med Rehabil* 1997 Feb;78(2):208-12.

Singh NA, Quine S, Clemson LM, et al. Effects of high-intensity progressive resistance training and targeted multidisciplinary treatment of frailty on mortality and nursing home admissions after hip fracture: a randomized controlled trial. *JAMDA* 2012;13:24-30.

Sylliaas H, Brovold T, Wyller TB, Bergland A. Progressive strength training in older patients after hip fracture: a randomized controlled trial. *Age Aging* 2011;40:221-227.

Yang Y, Komisar V, Shishov N, et al. The Effect of Fall Biomechanics on Risk for Hip Fracture in Older Adults: A Cohort Study of Video-Captured Falls in Long-Term Care. *J Bone Miner Res*. 2020;35(10):1914-1922. doi:10.1002/jbmr.4048

## **10. VIBRATION**

Beck BR, Norling TL. The effect of 8 mos of twice-weekly low-or higher intensity WBV on risk factors for postmenopausal hip fracture. *Am J Phys Med Rehabil* Dec 2010;89(12):997-1009.

Bernard B, Nelson N, Estill CF, Fine L. The NIOSH review of hand-arm vibration syndrome: vigilance is crucial. National Institute of Occupational Safety and Health. *Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine* Sep 1998;40(9):780-785.

Falcai MJ, Zamarioli A, Okubo R, de Paula FJ, Volpon JB. The osteogenic effects of swimming, jumping, and vibration on the protection of bone quality from disuse bone loss. *Scand J Med Sci Sports* 2014;Apr 30.

Kiel DP, Hannan MT, Barton BA, [Bouxsein ML](#), [Sisson E](#), [Lang T](#), [Allaire B](#), [Dewkett D](#), [Carroll D](#), [Magaziner J](#), [Shane E](#), [Leary ET](#), [Zimmerman S](#), [Rubin C](#). Low Magnitude Mechanical Stimulation to Improve Bone Density in Persons of Advanced Age: A Randomized, Placebo-Controlled Trial. *J Bone Miner Res* 2015;Jan 8.

Lai CL, Tseng SY, Chen CN, Liao WC, Wang CH, Lee MC Hsu PS. Effect of 6 months of whole body vibration on lumbar spine bone density in postmenopausal women: a randomized controlled trial. *Clin Interv Aging* 2013; 8:1603-1609.

Lam FM, Lau RW, Chung RC, Pang MY. The effect of whole body vibration on balance, mobility and falls in older adults: a systematic review and meta-analysis. *Maturitas* Jul 2012;72(3):206-213.

Lau RW, Liao LR, Yu F, Teo T, Chung RC, Pang MY. The effects of whole body vibration therapy on bone mineral density and leg muscle strength in older adults: a systematic review and meta-analysis. *Clinical Rehabilitation* Nov 2011;25(11):975-988.

Liphardt AM, Schipilow J, Hanley DA, Boyd S K. Bone quality in osteopenic postmenopausal women is not improved after 12 months of whole-body vibration training. *Osteoporos Int* 2015 26(3): 911-920.

Merriman HL, Braehler CJ, Jackson K. Systematically controlling for the influence of age, sex, hertz and time post-whole-body vibration exposure on four measures of physical performance in community-dwelling older adults: a randomized cross-over study. *Current Gerontology and Geriatrics Research* 2011;DOI:747094.

Merriman H, Jackson K. The effects of whole-body vibration training in aging adults: a systematic review. *J Geriatr Phys Ther* 2009;32(3):134-145.

Mikhael M, Orr R, Fiatarone Singh MA. The effect of whole body vibration exposure on muscle or bone morphology and function in older adults: a systematic review of the literature. *Maturitas* Jun 2010;66(2):150-157.

Muir J, Kiel DP, Rubin CT. Safety and severity of accelerations delivered from whole body vibration exercise devices to standing adults. *J Sci Med Sport* 2013;16(6):526-531.

Pollock RD, Martin FC, Newham DJ. Whole-body vibration in addition to strength and balance exercise for falls-related functional mobility of frail older adults: a single-blind randomized controlled trial. *Clinical Rehabilitation* Oct 2012;26(10):915-923.

Ruan XY, Jin FY, Liu YL, Peng ZL, Sun YG. Effects of vibration therapy on bone mineral density in postmenopausal women with osteoporosis. *Chinese Medical Journal* Jul 5 2008;121(13):1155-1158.

Rubin C, Recker R, Cullen D, Ryaby J, McCabe J, McLeod K. Prevention of postmenopausal bone loss by a low-magnitude, high-frequency mechanical stimuli: a clinical trial assessing compliance, efficacy, and safety. *J Bone Miner Res* Mar 2004;19(3):343-351.

Rubin C, Turner AS, Muller R, et al. Quantity and quality of trabecular bone in the femur are enhanced by a strongly anabolic, noninvasive mechanical intervention. *J Bone Miner Res* Feb 2002;17(2):349-357.

Slatkowska L, Alibhai SM, Beyene J, Cheung AM. Effect of whole-body vibration on BMD: a systematic review and meta-analysis. *Osteoporos Int* Dec 2010;21(12):1969-1980.

Slatkowska L, Alibhai SM, Beyene J, Hu H, Demaras A, Cheung AM. Effect of 12 months of whole-body vibration therapy on bone density and structure in postmenopausal women: a randomized trial. *Ann Intern Med* Nov 15 2011;155(10):668-679, W205.

Verschueren SM, Bogaerts A, Delecluse C, et al. The effects of whole-body vibration training and vitamin D supplementation on muscle strength, muscle mass, and bone density in institutionalized elderly women: a 6-month randomized, controlled trial. *J Bone Miner Res* Jan 2011;26(1):42-49.

von Stengel S, Kemmler W, Engelke K, Kalender WA. Effects of whole body vibration on bone mineral density and falls: results of the randomized controlled ELVIS study with postmenopausal women. *Osteoporos Int* Jan 2011;22(1):317-325.

Wysocki A, Butler M, Shamliyan T, Kane RL. Whole-body vibration therapy for osteoporosis: state of the science. *Ann Intern Med* Nov 15 2011;155(10):680-686, W206-613.

Zha DS, Zhu QA, Pei WW, et al. Does whole-body vibration with alternative tilting increase bone mineral density and change bone metabolism in senior people? *Aging Clin Exp Res* Feb 2012;24(1):28-36.

## **11. MEN**

Bolam K, von Uffelen J, Taaffe D. The Effect of Physical Exercise on Bone Density in Middle-aged and Older Men: a systematic review. *Osteoporos Int* 2013;24:2749-62.

Cauley JA, Harrison SL, Cawthon PM, Ensrud KE, Danielson ME, Orwoll E, Mackey DC. Objective measures of physical activity, fractures and falls: the osteoporotic fractures in men study. *J Am Geriatr Soc* 2013;61(7):1080-1088.

Harding AT, Weeks BK, Lambert C, Watson SL, Weis LJ, Beck BR. Effects of supervised high-intensity resistance and impact training or machine-based isometric training on regional bone geometry and strength in middle-aged and older men with low bone mass: The LIFTMOR-M semi-randomised controlled trial. *Bone*. 2020;136:115362. doi:10.1016/j.bone.2020.115362

Harding AT, Weeks BK, Lambert C, Watson SL, Weis LJ, Beck BR. A Comparison of Bone-Targeted Exercise Strategies to Reduce Fracture Risk in Middle-Aged and Older Men with Osteopenia and Osteoporosis: LIFTMOR-M Semi-Randomized Controlled Trial. *J Bone Miner Res*. 2020;35(8):1404-1414. doi:10.1002/jbmr.4008

Katzman WB, Miller-Martinez D, Marshall LM, Lane NE, Kado DM. Kyphosis and paraspinal muscle composition in older men: a cross-sectional study for the osteoporotic fractures in men (MrOS) research group. *BMC Musculoskelet Disord* 2014; 15(1):19.

Tveit M, Rosengren B, Nilsson J et al. Bone mass following physical activity in young years: a mean 39-year prospective controlled study in men. *Osteoporos Int* 2013;24:1389-97.

Warden SJ, Mantila-Roosa SM, Kersh ME, Hurd AL, Fleisig GS, Pandy MG, Fuchs RK. Physical activity when young provides lifelong benefits to cortical bone size and strength in men. *Proc Natl Acad Sci USA* 2014 111(14): 5337-5342.

## **12. CHILDREN/ADOLESCENTS/YOUNG ADULTS**

Baxter-Jones AD, Faulkner RA, Forwood M, Mirwald RL, Bailey DA. Bone mineral accrual from 8 to 30 years of age: an estimation of peak bone mass. *J. Bone Miner Res* 2011; 26(8):1729Y39.

Baxter-Jones AD, Kontulainen SA, Faulkner RA, Bailey DA. A longitudinal study of the relationship of physical activity to bone mineral accrual from adolescence to young adulthood. *Bone* 2008; 43:1101-7.

Blimkie CJ, Rice S, Webber CE, Martin J, Levy D, Gordon CL. Effects of resistance training on bone mineral content and density in adolescent females. *Can J Physiol Pharmacol* 1996;74:1025-1033.

Callreus M; McGuigan F; Ringsberg K; Akesson K. Self-reported Recreational Exercise Combining Regularity and Impact is Necessary to Maximize Bone Mineral Density in Young Adult Women. *Osteoporos Int* 2012;23:2517-26.

Duckham, R. L., Baxter-Jones, A. D., Johnston, J. D., Vatanparast, H., Cooper, D., & Kontulainen, S. Does physical activity in adolescence have site-specific and sex-specific benefits on young adult bone size, content, and estimated strength? *J Bone Miner Res* 2014;29(2), 479-486.

Faigenbaum, AD and Meyer, GD. Pediatric Resistance Training: Benefits, Concerns, and Program Design Considerations. *Current Sports Medicine Reports* 2010; 9:161-168.

Faigenbaum, AD and Myer, GD. Resistance training amount young athletes: safety, efficacy and injury prevention effects. *Br J Sports Med* 2010; 44 (1): 56-63.

Ferry B, Lespessailles E, Rochcongar P, Duclos M, Courteix D. Bone health during late adolescence: Effects of an 8-month training program on bone geometry in female athletes. *Joint Bone Spine* 2012 Mar 9.

Fuchs RK, Bauer JJ, Snow CM. Jumping improves hip and lumbar spine bone mass in prepubescent children: a randomized controlled trial. *J Bone Miner Res*. 2001; 16:148-56.

Gunter K, Baxter-Jones AD, Mirwald RL, et al. Impact exercise increases BMC during growth: an 8year longitudinal study. *J. Bone Miner Res* 2008; 23:986-93.

Gunter KB, Almstedt HC, Janz KF. Physical Activity in Childhood May Be the Key to Optimizing Lifespan Skeletal Health. *Exerc Sport Sci Rev* 2012, 40:13-31.

Heinonen A, Sievanen H, Kannus P, Oja P, Pasanen M, Vuori I. High impact exercise and bones of growing girls: a 9-month controlled trial. *Osteoporos Int* 2000; 11:1010-17.

Hind K, Burrows M. Weight-bearing exercise and bone mineral accrual in children and adolescents: A review of controlled trials. *Bone* 2007;40(1):14–27.

Jackowski SA, Faulkner RA, Farthing JP, Kontulainen SA, Beck TJ, Baxter-Jones AD. Peak lean tissue mass accrual precedes changes in bone strength indices at the proximal femur during the pubertal growth spurt. *Bone* 2009; 44(6):1186–1190.

Janz KF, Burns TL, Levy SM, et al. Everyday activity predicts bone geometry in children: The Iowa Bone Development Study. *Med Sci Sports Exerc.* 2004;36(7):1124–31.

Janz KF, Gilmore JME, Levy SM, Letuchy EM, Burns TL, Beck TJ. Physical activity and femoral neck bone strength during childhood: the Iowa Bone Development Study. *Bone* 2007; 41:216-22.

Janz KF, Letuchy EM, Eichenberger Gilmore JM, et al. Early physical activity provides sustained bone health benefits later in childhood. *Med Sci Sports Exerc* 2010; 42:1072-8.

Macdonald HM, Kontulainen SA, Khan KM, McKay HA: Is a school-based physical activity intervention effective for increasing tibial bone strength in boys and girls? *J Bone Miner Res* 2007, 22:434-446.

MacKelvie KJ, Khan KM, Petit MA, Janssen PA, McKay HA. A school-based exercise intervention elicits substantial bone health benefits: a 2-year randomized controlled trial in girls. *Pediatrics* 2003; 112(6 Pt 1):e447.

MacKelvie KJ, Petit MA, Khan KM, Beck TJ, McKay HA. Bone mass and structure are enhanced following a 2-year randomized controlled trial of exercise in prepubertal boys. *Bone* 2004; 34(4):755-64.

McKay, HA, McLean L, Petit, M MacKelvie-O’Briedn, K, Janssen, P, Beck, T and KM Khan. "Bounce at the Bell": a novel program of short bouts of exercise improves proximal femur bone mass in early pubertal children. *Br J Sports Med* 2005 39: 521-526

Mosti MP, Carlsen T, Aas E, Hoff J, Stunes AK, & Syversen U. Maximal strength training improves bone mineral density and neuromuscular performance in young adult women. *J Strength Cond Res* 2014;27(10): 2879-2886.

Nichols DL, Sanborn CF, Love AM. Resistance training and bone mineral density in adolescent females. *J Pediatr* 2001;139:494-500.

Petit MA, McKay HA, MacKelvie KJ, Heinonen A, Khan KM, Beck TJ. A randomized school-based jumping intervention confers site and maturity-specific benefits on bone structural properties in girls: a hip structural analysis study. *J Bone Miner Res.* 2002; 17:363Y72.

Ratel, S. High-intensity and Resistance Training and Elite Young Athletes. *Med Sport Sci* 2011. 56: 84-96.

Tenforde, AS and Fredericson, M. Influence of Sports Participation on Bone Health in the Young Athlete: A Review of the Literature. *Phys Med and Rehabil* 2011; 3:161-8.

Tveit M, Rosengren BE, Nilsson JA, & Karlsson MK. Exercise in youth: High bone mass, large bone size, and low fracture risk in old age. *Scand J Med Sci Sports* Aug 2014.

Warden SJ, Davis IS, Fredericson M. Management and prevention of bone stress injuries in long-distance runners. *J Orthop Sports Phys Ther* 2014;44(10): 749-765.

Weeks BK, Young CM, Beck BR. Eight months of regular in-school jumping improves indices of bone strength in adolescent boys and Girls: the POWER PE study. *J Bone Miner Res* 2008; 23:1002-11.

Wiebe PN, Blimkie CJ, Farpour-Lambert N, et al. Effects of single-leg drop-landing exercise from different heights on skeletal adaptations in prepubertal girls: a randomized controlled study. *Pediatr Exerc Sci* 2008;20:211-28.

Witzke KA, Snow CM. Effects of plyometric jump training on bone mass in adolescent girls. *Med Sci Sports Exerc* 2000; 32:1051-7.